DOI: 10.1111/1365-2656.13698

CONCEPT

Journal of Animal Ecology

Animal tracking moves community ecology: Opportunities and challenges

Raul Costa-Pereira¹ Remington J. Moll² Brett R. Jesmer^{3,4,5} Kernet Valter Jetz^{4,5}

¹Departamento de Biologia Animal, Instituto de Biociências, Universidade Estadual de Campinas, Campinas, Brazil

²Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH, USA

³Department of Fish and Wildlife Conservation, Virginia Tech, Blacksburg, VA, USA

⁴Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA

⁵Center for Biodiversity and Global Change, Yale University, New Haven, CT, USA

Correspondence Walter Jetz Email: walter.jetz@yale.edu

Funding information

Fundação de Amparo à Pesquisa do Estado de São Paulo, Grant/Award Number: 2020/11953-2; Instituto Serrapilheira, Grant/Award Number: Serra -R-2011-37572: National Aeronautics and Space Administration, Grant/Award Number: 80NSSC18K; Yale University, Max Planck - Yale Center for Biodiversity Movement and Global Change

Handling Editor: Allert Imre Bijleveld

Abstract

- 1. Individual decisions regarding how, why and when organisms interact with one another and with their environment scale up to shape patterns and processes in communities. Recent evidence has firmly established the prevalence of intraspecific variation in nature and its relevance in community ecology, yet challenges associated with collecting data on large numbers of individual conspecifics and heterospecifics have hampered integration of individual variation into community ecology.
- 2. Nevertheless, recent technological and statistical advances in GPS-tracking, remote sensing and behavioural ecology offer a toolbox for integrating intraspecific variation into community processes. More than simply describing where organisms go, movement data provide unique information about interactions and environmental associations from which a true individual-to-community framework can be built.
- 3. By linking the movement paths of both conspecifics and heterospecifics with environmental data, ecologists can now simultaneously quantify intraspecific and interspecific variation regarding the Eltonian (biotic interactions) and Grinnellian (environmental conditions) factors underpinning community assemblage and dynamics, yet substantial logistical and analytical challenges must be addressed for these approaches to realize their full potential.
- 4. Across communities, empirical integration of Eltonian and Grinnellian factors can support conservation applications and reveal metacommunity dynamics via tracking-based dispersal data. As the logistical and analytical challenges associated with multi-species tracking are surmounted, we envision a future where individual movements and their ecological and environmental signatures will bring resolution to many enduring issues in community ecology.

KEYWORDS

dispersal, ecological interactions, environmental niche, GPS-tracking, intraspecific variation, remote sensing

Raul Costa-Pereira, Remington J. Moll and Brett R. Jesmer contributed equally.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. Journal of Animal Ecology published by John Wiley & Sons Ltd on behalf of British Ecological Society.

1 | UNPACKING COMMUNITY PROCESSES

Ecologists have long sought to understand the forces governing the structure and functioning of the multilayered fabric of life, which involves recognizing a hierarchy of processes operating from individuals to communities (Levin, 1992). The inherent complexity of ecological systems has forced ecologists to balance realism and tractability in their models, which has resulted in a focus on species rather than individuals. Indeed, this widespread tendency is well justified given that interspecific differences are generally more conspicuous than intraspecific differences (Coulson, 2020). Nonetheless, the dynamics of animal communities are ultimately the products of individual decisions regarding how, why and when organisms move and interact with one another and with their environment (Potts et al., 2014; Schlägel et al., 2020; Spiegel et al., 2017). For instance, interspecific niche partitioning, a long-recognized condition favouring species coexistence (MacArthur & Levins, 1967), emerges from variation in resource use within and between species (Costa-Pereira, Araújo, et al., 2019). Both intraspecific and interspecific dietary niche differences are shaped by how conspecifics and heterospecifics navigate the trophic landscapes (e.g. by selecting food patches) (Pansu et al., 2019).

Although species-level studies have greatly advanced our understanding about community assembly and functioning over the last half century, a rapidly growing body of literature has revealed the significance of intraspecific ecological variation in communities (Bolnick et al., 2011). Generalist populations are often composed of groups of individuals that are resource or habitat specialists (Ingram et al., 2018; Schirmer et al., 2020) and this intraspecific variation can have stronger ecological effects than interspecific variation (Des Roches et al., 2018: Harrison et al., 2019). Therefore, individual ecological diversity plays a key role in shaping dynamics at the level of populations, communities and ecosystems (Allgeier et al., 2020; Bolnick et al., 2011; Ingram et al., 2018; Schirmer et al., 2020). Individuals within communities differ in their traits (Violle et al., 2012), trophic niches (Costa-Pereira, Araújo, et al., 2019), behaviours (Dantzer & Rubenstein, 2017; Harrison et al., 2019) and environmental associations (Carlson et al., 2021). Despite the importance of intraspecific and interspecific interactions and environmental associations in structuring communities (Thompson et al., 2020), understanding how these individual-level processes scale up to shape community dynamics remains an outstanding gap in knowledge.

The empirical development of an individual-to-community approach has been hindered by two major challenges. First, because of the enormous logistical challenge of collecting data at an individual level at high spatiotemporal resolutions, there have been historical empirical limitations of incorporating individual variation in a multispecies framework (Coulson, 2020). Second, studies have often focused on either interactions between organisms or environmental associations rather than both simultaneously (Soberón, 2007), creating conceptual and logistical discontinuities. We contend that movement data collected within a holistic conceptual framework and integrated across individuals and species has the potential to bring resolution to many enduring issues in community ecology.

2 | LINKING INDIVIDUAL MOVEMENTS TO COMMUNITY PROCESSES

Recent advances in GPS-tracking, biologging technologies (e.g. animal-borne video, proximity loggers, PIT-tags) and fine-scale remotely sensed data now enable simultaneous quantification of how individuals interact with conspecifics, heterospecifics and their environment (Nathan et al., 2022; Tuia et al., 2022). These technological advances have facilitated tracking greater numbers of individuals across multiple co-occurring species (e.g. Davidson et al., 2020; Johnson et al., 2013; Raymond et al., 2015) in both terrestrial and aquatic systems. The high spatiotemporal resolution of these data facilitates detailed quantification of interactions over long timescales-even lifetimes (Nathan et al., 2022). These 'quantitative biographies' have already yielded numerous new insights into the social behaviour of animals (King et al., 2018; Strandburg-Peshkin et al., 2015; Strandburg-Peshkin et al., 2017). However, these recent developments have rarely been applied in community-level contexts (Bro-Jørgensen et al., 2019) (see Section 9).

Tracking of individuals in a multi-species framework (Figure 1) has the potential to reveal key processes underpinning communities (Bro-Jørgensen et al., 2019; Milles et al., 2020; Schlägel et al., 2020). First, movement data facilitate more than just mapping biotic interactions at fine spatiotemporal scales. When coupled with high-resolution environmental data, individual movements also describe fine-scale variation in environmental associations both within and between species, providing fine-scale data to investigate how environmental filters drive community patterns (Bastille-Rousseau & Wittemyer, 2019). At broader scales, movement data can help elucidate metacommunity and regional diversity dynamics by providing a window through which we can observe dispersal and migration and their consequences on both source and receiver communities (Bauer & Hoye, 2014; Jesmer et al., 2018; Schlägel et al., 2020).

3 | USING MOVEMENT DATA TO INTEGRATE ELTONIAN AND GRINNELLIAN COMMUNITY PROCESSES

Interactions between conspecifics, heterospecifics and the environment form the backbone of community ecology. Understanding general patterns of biodiversity should therefore involve uncovering both the impacts and responses of an organism in relation to other organisms (i.e. biotic interactions)—referred to as Eltonian factors (Chase & Leibold, 2003; Soberón, 2007)—and the environmental conditions necessary to sustain neutral or positive fitness values (i.e. environmental associations)—referred to as Grinnellian factors (Thompson et al., 2020). Although ongoing work seeks to integrate these concepts under a unified theory of ecological niches (Chase & Leibold, 2003; Gravel et al., 2019; Peterson et al., 2011; Potts et al., 2014), Eltonian and Grinnellian factors have traditionally been studied in isolation. This is due to a paucity of data capable of supporting such an integration (i.e. data for interacting organisms collected



FIGURE 1 Eltonian and Grinnelian dynamics inferred from multi-species tracking data. *Panel (a)*: Tracks of five individuals from three different species reveal intraspecific and interspecific interactions through time, thereby enabling the construction of interaction topologies including both conspecifics and heterospecifics at an individual level. *Panel (b)*: Interactions in the Eltonian arena can be mapped via temporally explicit tracks, allowing for spatiotemporal analysis of interactions across landscapes. Therefore, the intersection of tracks with environmental data (e.g. remote sensing layers) in space and time quantifies environmental associations and facilitates assessments of population- and community-wide Grinnellian niche partitioning

over consistent spatial and temporal scales) (Soberón, 2007). By linking modern remote-sensing data with auxiliary biologging technologies (e.g. proximity loggers, animal-borne video), the tracking of individual organisms in a multi-species framework may now overcome these historical challenges and support synthesis. Here we offer a framework for leveraging these new technology-driven opportunities in community ecology and explore the key challenges that must be overcome to make this vision a reality.

4 | ELTONIAN FACTORS CAPTURED WITH TRACKING DATA

Elton defined the niche as the role a species plays in a given community, particularly 'its relations to food and enemies' (MacArthur & Levins, 1967). Contemporary theory has operationalized this concept by focusing on the impacts one species has on other organisms (Chase & Leibold, 2003; Letten et al., 2017). The influence of biotic interactions that shape this niche space, such as predation and competition, can be inferred probabilistically from analysis of individual movement data which is both spatially and temporally explicit (Isbell et al., 2018; Milner et al., 2021; Nathan et al., 2022; Schlägel et al., 2019; Suraci et al., 2022; Villegas-Ríos et al., 2020). These Eltonian factors vary by magnitude, frequency and type, thereby encompassing interactions often described at the species level in classical theory (Wootton, 1994). Importantly, movement data facilitate insight both into intraspecific (King et al., 2018; Strandburg-Peshkin et al., 2015) and interspecific biotic interactions (Montiglio et al., 2019), thereby enabling the construction of community topologies (e.g. food webs) using individuals rather than species as functional units (Figure 1).

Classical work on Eltonian factors for species pairs is built upon consumer-resource models (Murdoch et al., 2003) and has since been expanded to include non-consumptive factors, whereby phenotypic changes mitigate negative interactions (e.g. anti-predator behaviours; Abrams, 1995). Sequential and simultaneous location data from multiple moving organisms (both conspecifics and heterospecifics) enable inference regarding Eltonian factors from patterns of attraction, repulsion or neutrality (Milner et al., 2021; Potts et al., 2014; Schlägel et al., 2019). For example, direct predation is inferred via cessation of motion or clustered locations surrounding kill sites (Anderson & Lindzey, 2003). When combined with auxiliary data (e.g. diet composition) and other biologging measures (e.g. accelerometers, heart rate loggers) (Williams et al., 2020), individual movements can reveal how consumptive and non-consumptive factors combine to shape fitness-optimizing behaviour. For instance, a proximate threat of predator encounter can cause prey to move to less nutritious, but safer, foraging patches (Barnier et al., 2014) and

lead to trophic cascades (Ford et al., 2014). Certain Eltonian factors are linked to symmetric spatiotemporal associations (e.g. positivepositive as in cooperative foraging), while others result in asymmetries (e.g. negative-positive, as in prey fleeing a cursorial predator) (Villegas-Ríos et al., 2020). Such asymmetries underscore the need for careful analyses to disentangle pattern from process (Freilich et al., 2018).

Movement data can also reveal indirect Eltonian factors, which occur when the effects of one species on another are mediated by a third species (e.g. apparent competition). Like their direct counterparts, indirect factors propagate along consumptive and nonconsumptive pathways (Abrams, 1995; Gil et al., 2018). Importantly, these factors have analogues on the individual organizational level, although studies at this resolution remain rare. For example, multispecies tracking studies on carnivores and ungulates have revealed complex intraguild interspecific avoidance based upon dominance hierarchies and associated mechanisms of community niche partitioning (Dröge et al., 2017; Vanak et al., 2013). However, little understanding of how individual variation influences such community dynamics exists (Gil et al., 2018). Multi-species movement studies represent an exciting new research avenue because the mechanistic underpinnings of complex community processes have likely been masked by aggregative spatiotemporal patterns at the population level (Bolnick et al., 2011; Peterson et al., 2020; Potts et al., 2014).

5 | GRINNELLIAN FACTORS CAPTURED WITH TRACKING DATA

Since Grinnell's original conceptualization, a species' environmental niche has been defined as the suite of conditions necessary to sustain viable populations (Chase & Leibold, 2003). Ecologists' ability to quantify such Grinnellian factors is rapidly developing as advances in sensing technologies, data processing and computational modelling have led to a growing number of remotely sensed products that capture environmental conditions at increasingly fine spectral, temporal and spatial resolution at a near-global extent (Anderson, 2018; He et al., 2015; Jetz et al., 2016; Mertes et al., 2020; Randin et al., 2020; Tuia et al., 2022; Wilson & Jetz, 2016). Furthermore, airborne sensing (e.g. LiDAR) offers even higher spectral and spatiotemporal resolution and detailed habitat characterizations (Asner et al., 2017; Carrasco et al., 2019). Simultaneously, tracking technologies have become more miniaturized and efficient, thereby facilitating tracking of smaller-bodied species and finer-scale temporal sampling of animal movement (Kays et al., 2015; Wikelski et al., 2007). Intersecting multi-species movement tracks with remotely sensed data allows quantifying Grinnellian factors for conspecifics and heterospecifics in *n*-dimensional environmental niche space (Carlson et al., 2021) (Figure 1). In turn, this niche space can be projected into geographical space to map distributional areas (Colwell & Rangel, 2009), allowing ecologists to assess whether and how intraspecific and interspecific variation in environmental niche space affects community dynamics (Figure 1).

Fundamental to quantifying Grinnellian factors is an understanding of the fitness consequences of environmental conditions (Pulliam, 2000). Although the many data types (e.g. presence only, presence-absence survey data, expert range maps) used to determine Grinnellian niches require ancillary studies to quantify individual fitness, tracking data can simultaneously assess fitness and the environmental conditions experienced by individuals from multiple co-occurring species. For example, individual movement patterns enable ecologists to remotely evaluate survival, parturition and recruitment across heterogeneous landscape conditions (DeMars et al., 2013; Hooven et al., 2022). In this way, spatiotemporally explicit movement data permit identifying when and where individuals, populations and species experience favourable conditions.

6 | TOWARDS AN INTEGRATED UNDERSTANDING

Movement data enable joint assessment, and thus integration, of Eltonian and Grinnellian factors at the level of individuals, populations and communities (Figure 1). This integration allows ecologists to unify niche concepts and may illuminate how intraspecific niche variation influences classical paradigms in community ecology (e.g. species coexistence) (Bolnick et al., 2011; Milles et al., 2020). To successfully integrate across niche concepts and organismal scales, movement models need to jointly estimate (a) the impacts individuals have on each other (both conspecifics and heterospecifics) and (b) the conditions individuals need to sustain positive fitness values. Such joint quantifications are enabled by a growing set of methods and tools, particularly those from social behavioural research (Bro-Jørgensen et al., 2019; King et al., 2018; Nathan et al., 2022; Strandburg-Peshkin et al., 2015; Tuia et al., 2022), but will require judgement around balancing analytical complexity with logistical and interpretational tractability (see Section 9).

The integrative potential of movement data for community ecology is further enhanced by complementary sources of ecological data. For instance, when combined with movement data, a diverse array of methods (e.g. stable isotopes, DNA metabarcoding, video and proximity-enabled collars) now support the characterization of intraspecific and interspecific niche partitioning and plasticity. Such auxiliary information enables deeper insights into the behavioural mechanisms shaping niches (Bastille-Rousseau & Wittemyer, 2019). For instance, video and proximity-enabled tracking devices supply fine-scale behavioural data on interspecific interactions that may not be fully captured by locational data alone (e.g. physical contact between individuals) (Lavelle et al., 2012; Moll et al., 2007). Likewise, combining animal tracking with stable isotope or DNA metabarcoding data can help uncover how foraging decisions may influence patterns of movement and space use (Atkins et al., 2019; Bradshaw et al., 2017; Votier et al., 2010). To date, studies of such sophistication have predominantly focused on individuals of a single species (see Section 9), but multi-species extensions represent an exciting avenue for future research.

7 | FROM SINGLE TO MULTIPLE COMMUNITIES AND ECOSYSTEMS

The ability of movement data to quantify processes shaping community dynamics extends beyond single communities. As Eltonian and Grinnellian factors vary along environmental gradients (Gravel et al., 2019), tracking individuals from sets of species across multiple communities will enlighten how local contexts (e.g. local diversity, environmental heterogeneity) affect individual interactions and thus species niches (Figure 2). For instance, comparing how interactions between conspecifics and heterospecifics change along temperature gradients or with the presence of predators are, respectively, promising research agendas for understanding how climate change and defaunation impact the organization of communities (Barnier et al., 2014; Veldhuis et al., 2020).

Community ecology has experienced a change in its focus from single to multiple, connected communities. Metacommunity theory proposes that the interplay of within- and among-community factors drives regional biodiversity patterns (Leibold & Chase, 2017; Thompson et al., 2020). In this context, individual movements can provide critical information about among-community connectivity via biotic links. Dispersal is the key process governing spatial dynamics across metacommunities (Leibold & Chase, 2017; Thompson et al., 2020); therefore, tracking data can answer open questions about the causes of individual dispersal and their implications for populations and communities (Schlägel et al., 2020). For instance, mapping pre-dispersal interactions and environmental associations may reveal factors triggering dispersal by specific individuals within populations, as well as how dispersal impacts both recipient and source metacommunities (Figure 2). Finally, individual movements can also help understand links among ecosystems, sometimes separated by thousands of kilometres (Alerstam et al., 2003; Bauer & Hoye, 2014). Because animals move substantial amounts of matter and energy across ecosystem boundaries (Schmitz et al., 2018), movement data can uncover the magnitude of spatial coupling between ecosystems, allowing the development of models capable of predicting the impacts of disruptions in meta-ecosystem fluxes due to anthropogenic barriers to movement.

8 | FROM BASIC TO APPLIED COMMUNITY ECOLOGY

In a rapidly changing world, the combination of animal movement and remote sensing data offers powerful conservation perspectives for safeguarding biodiversity (Nathan et al., 2022; Tuia et al., 2022). For example, Grinnellian niche assessments support myriad uses for modelling fine-scale species distributions and animal-habitat relationships. Such assessments are especially effective for herbivores because remote sensing data can often directly capture their relevant niche factors (e.g. NDVI), although matching remote sensing product resolution with fine-scale tracking data remains an ongoing challenge, especially for predators (Suraci et al., 2022). The output of such models can help predict the current and future composition of communities, which, in turn, can strategically guide conservation action (e.g. identify diversity hotspots) (Hazen et al., 2013). For instance, by tracking individuals from several (n = 6-21) species of birds and mammals inhabiting the same region at the same time, specific marine regions critical for breeding and foraging at the community level were identified (Davies et al., 2021; Raymond et al., 2015). These emerging patterns from individual-level data in a multi-species context are vital for informing policy regarding the establishment of protected areas.

Individual movement data also have the potential to support mechanistic and spatiotemporally explicit predictions of how anthropogenic impacts (e.g. land use change, species introductions) will change interactions between conspecifics and heterospecifics (Kays et al., 2015; Nathan et al., 2022; Veldhuis et al., 2020). For example, while it is widely recognized that wildlife often responds to human activity by becoming more nocturnal or limiting movements (Gaynor et al., 2018; Tucker et al., 2018), individual-level analyses could reveal how responses to human activity vary within and across species. In turn, individual behavioural responses are likely to alter the dynamics of populations and communities (Bolnick et al., 2011; Laskowski et al., 2022) and offer insights regarding the behavioural phenotypes and behavioural syndromes most likely to persist under scenarios of increased anthropogenic disturbance (Schell et al., 2021).

Accounting for intraspecific variation has become increasingly important for successful conservation (Des Roches et al., 2021; Merrick & Koprowski, 2017). Although management plans often use 'average' (i.e. species level) behaviour and diet data to support decisions, individual variation in Grinnellian and Eltonian factors may be linked to particular phenotypes that disproportionately contribute to a population's dynamics or persistence (Costa-Pereira, Toscano, et al., 2019; Durell, 2000; Montgomery et al., 2018). Work on the individuality of habitat selection and movement has found that intraspecific variation can be stronger than that of even interspecific differences (Harrison et al., 2019; Montgomery et al., 2018). Indeed, considering animal individuality in conservation efforts is emerging as an effective solution for mitigating human-wildlife conflict, which is often driven by particular phenotypes (e.g. bold individuals) (Barrett et al., 2019; Honda et al., 2018). The inherent link between individual movements and community dynamics holds an increasingly important role in conservation planning and management, thereby potentially advancing our ability to effectively address the global biodiversity crisis (Kays et al., 2015; Merrick & Koprowski, 2017; Nathan et al., 2022).

9 | CHALLENGES

Multi-species tracking datasets are becoming more commonplace (Nathan et al., 2022; Wilmers et al., 2015), yet financial, technical



FIGURE 2 Inferring Eltonian and Grinnelian dynamics from single to multiple communities. Panels depict how communities vary by the number of species present, the number of individuals per species, the movement behaviour of such individuals and local environmental characteristics. Community 1 harbours three tracked species, while in community 2, species 2 experiences competition release and in community 3 species 1 and 2 experience predation release. The dispersal event (individual_e) depicted between communities 1 and 2 emphasizes the power of tracking data to quantify biotic links across communities

and operational challenges must be overcome if quantifying Eltonian and Grinnellian factors from movement data is to become conventional practice. First, these opportunities are currently limited to species capable of carrying tracking devices. New technological developments (e.g. icarus.mpg.de), however, are poised to expand the catalogue of species and guilds possessing movement data (Jetz et al., 2022; animallives.org). Importantly, some ecological actors will inevitably be missed since it is impossible to track every individual in a community, and these missed actors could induce substantial bias in community-level inference. For example, Creel et al. (2013) found that failing to account for missed interactions between GPS-collared predators and prey could underestimate antipredator behavioural responses by an order of magnitude or more. This challenge can be further intensified when interacting species exhibit movement patterns across disparate spatial scales (Suraci et al., 2022). For instance, a recent study of wide-ranging wolves *Canis lupus* and more sedentary elk *Cervus elaphus* recorded more than 36,000 and 13,000 GPS locations of these species, respectively, but documented only 453 'encounters' where conspecifics were co-located within 1,000 m of each other (Cusack et al., 2020). Addressing these challenges will be system specific and likely involve combining experimental designs that minimize sampling bias with probabilistic modelling (Farage et al., 2021; Gupte et al., 2022; Schlägel et al., 2019), individual-based simulation (King et al., 2018) and auxiliary data to 'fill-in' critical gaps for unsampled individuals (e.g. camera traps and video-enabled tracking devices capable of capturing interspecific and intraspecific interactions with unmarked individuals).

Although such auxiliary data provide contextual biological information that compliment movement data, there is still a critical challenge of matching spatiotemporal resolution of multiple data types, which can be quite disparate (e.g. minute-level resolution of movement data and season-level resolution of stable isotope data). This challenge is not trivial, but new data fusion algorithms are rapidly developing (Brum-Bastos et al., 2020; Gupte et al., 2022; Marvin et al., 2016). We expect such methods to constitute an important area of growth moving forward. Furthermore, because modern GPS-tracking devices can capture data at temporal resolutions of minutes or less, tracking individuals from multiple species over appreciable ecological timescales causes the number of data points soar into the tens of millions. Recent single-species work on social group dynamics, however, has laid the methodological and analytical foundations from which massive amounts of individual level-data can be processed, analysed and interpreted (King et al., 2018; Silk et al., 2018; Strandburg-Peshkin et al., 2015). Thus, the foundation for managing this volume of the data as well as the interpretability and validation of results has been laid and additional, and novel methods that are crucial for operationalizing an individual-to-community framework are developing rapidly (Gupte et al., 2022; Nathan et al., 2022; Noonan et al., 2021; Tuia et al., 2022).

The benefit of multispecies tracking datasets appears to be increasingly acknowledged and employed. For example, by tracking several species at the same time in same region, key breeding and foraging sites and migratory routes needed to sustain multiple species at a single site can be identified (Davies et al., 2021; Kauffman et al., 2021; Lowther et al., 2015). However, the tracking of multiple species in the same location is often conducted in isolation by different research groups and the data are typically not well integrated (Nathan et al., 2022). Such scenarios likely stem from differing research agendas and the substantial financial and logistical challenges associated with deploying large numbers of tracking devices on multiple co-occurring species, thereby highlighting the need for collaboration and data sharing if we are to move towards a more community-focused use of movement data (Davidson et al., 2020; Kays et al., 2022; Nathan et al., 2022; Urbano et al., 2021). Indeed, coordinating the field logistics alone for multi-species tracking studies comprises a major challenge that will likely only be overcome through increased coordination among research teams and an inclusive, transparent and collaborative approach to data collection and sharing.

10 | CONCLUDING REMARKS AND FUTURE VISION

The historical duality between Eltonian and Grinnellian approaches in community ecology has hampered our understanding of how communities are organized in space and time (Soberón, 2007). Additionally, despite growing evidence of the critical role of individual variation in community-level processes (Bolnick et al., 2011; Costa-Pereira, Araújo, et al., 2019; Milles et al., 2020), empirical studies are still largely species oriented. These shortcomings stem in part from our inability to simultaneously quantify Eltonian and Grinnellian factors at the individual level in a multi-species context. We hope the framework presented here will help ecologists overcome these shortcomings by simultaneously capturing interactions between organisms (conspecifics and heterospecifics) and their environment.

This individual-level framework outlined here should complement—and needs to be complemented by—more traditional population- and community-level approaches to estimate key parameters such as population size, phenotypic trait distribution and local community structure. Combining these data sources certainly entails overcoming major logistical, financial and analytical challenges, yet we are optimistic that recent and future technical developments are beginning to coalesce into Hutchinson's dream toolbox: a toolset capable of simultaneously quantifying multiple niche dimensions across multiple coexisting species. The intuitive, but still largely unexplored, integration between animal movements and community ecology has the potential to shed light on long-standing questions in ecological theory and help develop new predictive models for gauging the effects of global change on communities.

ACKNOWLEDGEMENTS

This study resulted from a symposium on 'Linking Individual Behavior to Community Responses in Changing Landscapes' held by the Max Planck-Yale Center for Biodiversity Movement and Global Change (MPYC, mpyc.yale.edu) at Yale University 13-14 May 2019. We would like to thank the participants for discussions that inspired this work, namely Ben Carlson, Blair Costello, Meg Crofoot, Diego Ellis Soto, Vanessa Ezenwa, Damien Farine, Richard Li, Roland Kavs, Doris Klein, Tyler Kartzinel, Katherine Mertes, Mathew Mutinda, Dan Rubenstein, Dustin Rubenstein, Ryan Pavlick, Jared Stabach, Erica Stuber, Ruth Oliver and Martin Wikelski. We thank D. Farine, B. Carlson, T. Kartzinel, M. Wikelski, G. Passoni and the Associate editor for valuable comments that helped us to improve our manuscript. W.J. acknowledges Yale University support for the MPYC symposium and National Aeronautics and Space Administration (NASA) grant 80NSSC18K. R.C.-P. is supported by the Instituto Serrapilheira (grant number Serra-R-2011-37572) and grant #2020/11953-2, São Paulo Research Foundation (FAPESP).

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

AUTHORS' CONTRIBUTIONS

All authors conceived the ideas, wrote collectively the manuscript and contributed critically to the drafts. We all gave final approval for publication.

DATA AVAILABILITY STATEMENT

Data will not be archived because this manuscript does not use any data.

ORCID

Raul Costa-Pereira https://orcid.org/0000-0003-2370-5866 Brett R. Jesmer https://orcid.org/0000-0002-2868-7161

REFERENCES

- Abrams, P. A. (1995). Implications of dynamically variable traits for identifying, classifying, and measuring direct and indirect effects in ecological communities. *The American Naturalist*, 146, 112–134.
- Alerstam, T., Hedenström, A., & Åkesson, S. (2003). Long-distance migration: Evolution and determinants. Oikos, 103, 247–260.
- Allgeier, J. E., Cline, T. J., Walsworth, T. E., Wathen, G., Layman, C. A., & Schindler, D. E. (2020). Individual behavior drives ecosystem function and the impacts of harvest. *Science Advances*, 6, eaax8329.
- Anderson, C. B. (2018). Biodiversity monitoring, earth observations and the ecology of scale. *Ecology Letters*, 21, 1572–1585.
- Anderson, C. R., & Lindzey, F. G. (2003). Estimating cougar predation rates from GPS location clusters. The Journal of Wildlife Management, 67, 307–316.
- Asner, G. P., Martin, R. E., Knapp, D. E., Tupayachi, R., Anderson, C. B., Sinca, F., Vaughn, N. R., & Llactayo, W. (2017). Airborne laserguided imaging spectroscopy to map forest trait diversity and guide conservation. *Science*, 355, 385–389.
- Atkins, J. L., Long, R. A., Pansu, J., Daskin, J. H., Potter, A. B., Stalmans, M. E., Tarnita, C. E., & Pringle, R. M. (2019). Cascading impacts of large-carnivore extirpation in an African ecosystem. *Science*, *364*, 173–177.
- Barnier, F., Valeix, M., Duncan, P., Chamaillé-Jammes, S., Barre, P., Loveridge, A. J., Macdonald, D. W., & Fritz, H. (2014). Diet quality in a wild grazer declines under the threat of an ambush predator. Proceedings of the Royal Society B: Biological Sciences, 281, 20140446.
- Barrett, L. P., Stanton, L. A., & Benson-Amram, S. (2019). The cognition of 'nuisance' species. Animal Behaviour, 147, 167–177.
- Bastille-Rousseau, G., & Wittemyer, G. (2019). Leveraging multidimensional heterogeneity in resource selection to define movement tactics of animals. *Ecology Letters*, 22, 1417–1427.
- Bauer, S., & Hoye, B. J. (2014). Migratory animals couple biodiversity and ecosystem functioning worldwide. *Science*, 344, 1242552.
- Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., Rudolf, V. H. W., Schreiber, S. J., Urban, M. C., & Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, *26*, 183–192.
- Bradshaw, P. J., Broderick, A. C., Carreras, C., Inger, R., Fuller, W., Snape, R., Stokes, K. L., & Godley, B. J. (2017). Satellite tracking and stable isotope analysis differential recruitment among foraging areas in green turtles. *Marine Ecology Progress Series*, 582, 201–214. https:// doi.org/10.3354/meps12297
- Bro-Jørgensen, J., Franks, D. W., & Meise, K. (2019). Linking behaviour to dynamics of populations and communities: Application of novel approaches in behavioural ecology to conservation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 374, 20190008.
- Brum-Bastos, V., Long, J., Church, K., Robson, G. G., de Paula, R., & Demšar, U. (2020). Multi-source data fusion of optical satellite imagery to characterize habitat selection from wildlife tracking data. *Ecological Informatics*, 60, 101149. https://doi.org/10.1016/j. ecoinf.2020.101149.
- Carlson, B. S., Rotics, S., Nathan, R., Wikelski, M., & Jetz, W. (2021). Individual environmental niches in mobile organisms. *Nature Communications*, 12, 4572.
- Carrasco, L., Giam, X., Papeş, M., & Sheldon, K. (2019). Metrics of Lidarderived 3D vegetation structure reveal contrasting effects of horizontal and vertical forest heterogeneity on bird species richness. *Remote Sensing*, 11(7), 743. https://doi.org/10.3390/rs11070743

- Chase, J. M., & Leibold, M. A. (2003). Ecological niches: Linking classical and contemporary approaches. University of Chicago Press.
- Colwell, R. K., & Rangel, T. F. (2009). Hutchinson's duality: The once and future niche. Proceedings of the National Academy of Sciences of the United States of America, 106, 19651–19658.
- Costa-Pereira, R., Araújo, M. S., Souza, F. L., & Ingram, T. (2019). Competition and resource breadth shape niche variation and overlap in multiple trophic dimensions. *Proceedings of the Royal Society B: Biological Sciences, 286,* 20190369.
- Costa-Pereira, R., Toscano, B., Souza, F. L., Ingram, T., & Araújo, M. S. (2019). Individual niche trajectories drive fitness variation. *Functional Ecology*, 33, 1734–1745.
- Coulson, T. (2020). Ecology and evolution is hindered by the lack of individual-based data. In A. Dobson, R. D. Holt, & D. Tilman (Eds.), Unsolved problems in ecology (pp. 107–114). Princeton University Press.
- Creel, S., Winnie, J. A., Jr., & Christianson, D. (2013). Underestimating the frequency, strength and cost of antipredator responses with data from GPS collars: An example with wolves and elk. *Ecology and Evolution*, *3*, 5189–5200.
- Cusack, J. J., Kohl, M. T., Metz, M. C., Coulson, T., Stahler, D. R., Smith, D. W., & MacNulty, D. R. (2020). Weak spatiotemporal response of prey to predation risk in a freely interacting system. *Journal of Animal Ecology*, 89, 120–131.
- Dantzer, B., & Rubenstein, D. R. (2017). Introduction to symposium: The developmental and proximate mechanisms causing individual variation in cooperative behavior. *Integrative and Comparative Biology*, 57, 560–565.
- Davidson, S. C., Bohrer, G., Gurarie, E., LaPoint, S., Mahoney, P. J., Boelman, N. T., Eitel, J. U. H., Prugh, L. R., Vierling, L. A., Jennewein, J., Grier, E., Couriot, O., Kelly, A. P., Meddens, A. J. H., Oliver, R. Y., Kays, R., Wikelski, M., Aarvak, T., Ackerman, J. T., ... Hebblewhite, M. (2020). Ecological insights from three decades of animal movement tracking across a changing Arctic. *Science*, *370*(6517), 712-715. https://doi.org/10.1126/science.abb7080
- Davies, T. E., Carneiro, A. P. B., Tarzia, M., Wakefield, E., Hennicke, J. C., Frederiksen, M., Hansen, E. S., Campos, B., Hazin, C., Lascelles, B., Anker-Nilssen, T., Arnardóttir, H., Barrett, R. T., Biscoito, M., Bollache, L., Boulinier, T., Catry, P., Ceia, F. R., Chastel, O., ... Dias, M. P. (2021). Multispecies tracking reveals a major seabird hotspot in the North Atlantic. *Conservation Letters*, 14(5). https://doi.org/10.1111/conl.12824
- DeMars, C. A., Auger-Méthé, M., Schlägel, U. E., & Boutin, S. (2013). Inferring parturition and neonate survival from movement patterns of female ungulates: A case study using woodland caribou. *Ecology* and Evolution, 3, 4149–4160.
- Des Roches, S., Pendleton, L. H., Shapiro, B., & Palkovacs, E. P. (2021). Conserving intraspecific variation for nature's contributions to people. *Nature Ecology & Evolution.*, 5, 574–582.
- Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., Schweitzer, J. A., & Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology & Evolution*, 2, 57–64.
- Dröge, E., Creel, S., Becker, M. S., & M'Soka, J. (2017). Spatial and temporal avoidance of risk within a large carnivore guild. *Ecology and Evolution*, 7, 189–199.
- Durell, S. E. A. L. V. D. (2000). Individual feeding specialisation in shorebirds: Population consequences and conservation implications. *Biological Reviews*, 75, 503–518.
- Farage, C., Edler, D., Eklöf, A., Rosvall, M., Pilosof, S., Farage, C., Edler, D., Eklöf, A., Rosvall, M., & Pilosof, S. (2021). Identifying flow modules in ecological networks using Infomap. *Methods in Ecology and Evolution*, 12(5), 778-786. https://doi. org/10.1111/2041-210x.13569
- Ford, A. T., Goheen, J. R., Otieno, T. O., Bidner, L., Isbell, L. A., Palmer, T. M., Ward, D., Woodroffe, R., & Pringle, R. M. (2014). Large

- Freilich, M. A., Wieters, E., Broitman, B. R., Marquet, P. A., & Navarrete, S. A. (2018). Species co-occurrence networks: Can they reveal trophic and non-trophic interactions in ecological communities? *Ecology*, 99, 690–699.
- Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence of human disturbance on wildlife nocturnality. *Science*, 360, 1232–1235.
- Gil, M. A., Hein, A. M., Spiegel, O., Baskett, M. L., & Sih, A. (2018). Social information links individual behavior to population and community dynamics. *Trends in Ecology & Evolution*, 33, 535–548.
- Gravel, D., Baiser, B., Dunne, J. A., Kopelke, J.-P., Martinez, N. D., Nyman, T., Poisot, T., Stouffer, D. B., Tylianakis, J. M., Wood, S. A., & Roslin, T. (2019). Bringing Elton and Grinnell together: A quantitative framework to represent the biogeography of ecological interaction networks. *Ecography*, 42(3), 401–415. https://doi.org/10.1111/ ecog.04006
- Gupte, P. R., Beardsworth, C. E., Spiegel, O., Lourie, E., Toledo, S., Nathan, R., & Bijleveld, A. I. (2022). A guide to pre-processing high-throughput animal tracking data. *Journal of Animal Ecology*, 91, 287–307.
- Harrison, P. M., Keeler, R. A., Robichaud, D., Mossop, B., Power, M., & Cooke, S. J. (2019). Individual differences exceed species differences in the movements of a river fish community. *Behavioral Ecology*, 30, 1289–1297.
- Hazen, E. L., Jorgensen, S., Rykaczewski, R. R., Bograd, S. J., Foley, D. G., Jonsen, I. D., Shaffer, S. A., Dunne, J. P., Costa, D. P., Crowder, L. B., & Block, B. A. (2013). Predicted habitat shifts of Pacific top predators in a changing climate. *Nature Climate Change*, *3*, 234–238.
- He, K. S., Bradley, B. A., Cord, A. F., Rocchini, D., Tuanmu, M.-N., Schmidtlein, S., Turner, W., Wegmann, M., & Pettorelli, N. (2015).
 Will remote sensing shape the next generation of species distribution models? *Remote Sensing in Ecology and Conservation*, 1(1), 4–18. https://doi.org/10.1002/rse2.7
- Honda, T., Iijima, H., Tsuboi, J., & Uchida, K. (2018). A review of urban wildlife management from the animal personality perspective: The case of urban deer. Science of the Total Environment, 644, 576–582.
- Hooven, N. D., Williams, K. E., Hast, J. T., McDermott, J. R., Crank, R. D., Jenkins, G., Springer, M. T., & Cox, J. J. (2022). Using low-fix rate GPS telemetry to expand estimates of ungulate reproductive success. Animal Biotelemetry, 10(1), 5. https://doi.org/10.1186/s4031 7-022-00276-0
- Ingram, T., Costa-Pereira, R., & Araújo, M. S. (2018). The dimensionality of individual niche variation. *Ecology*, *99*, 536–549.
- Isbell, L. A., Bidner, L. R., Van Cleave, E. K., Matsumoto-Oda, A., & Crofoot, M. C. (2018). GPS-identified vulnerabilities of savannahwoodland primates to leopard predation and their implications for early hominins. *Journal of Human Evolution*, 118, 1–13.
- Jesmer, B. R., Merkle, J. A., Goheen, J. R., Aikens, E. O., Beck, J. L., Courtemanch, A. B., Hurley, M. A., McWhirter, D. E., Miyasaki, H. M., Monteith, K. L., & Kauffman, M. J. (2018). Is ungulate migration culturally transmitted? Evidence of social learning from translocated animals. *Science*, *361*, 1023–1025.
- Jetz, W., Cavender-Bares, J., Pavlick, R., Schimel, D., Davis, F. W., Asner, G. P., Guralnick, R., Kattge, J., Latimer, A. M., Moorcroft, P., Schaepman, M. E., Schildhauer, M. P., Schneider, F. D., Schrodt, F., Stahl, U., & Ustin, S. L. (2016). Monitoring plant functional diversity from space. *Nature Plants*, 2, 16024.
- Jetz, W., Tertitski, G., Kays, R., Mueller, U., Wikelski, M., Åkesson, S., Anisimov, Y., Antonov, A., Arnold, W., Bairlein, F., Baltà, O., Baum, D., Beck, M., Belonovich, O., Belyaev, M., Berger, M., Berthold, P., Bittner, S., Blake, S., ... Zook, C. (2022). Biological Earth observation with animal sensors. *Trends in Ecology & Evolution*, 37(4), 293–298. https://doi.org/10.1016/j.tree.2021.11.011

- Johnson, H. E., Hebblewhite, M., Stephenson, T. R., German, D. W., Pierce, B. M., & Bleich, V. C. (2013). Evaluating apparent competition in limiting the recovery of an endangered ungulate. *Oecologia*, 171, 295–307.
- Kauffman, M. J., Cagnacci, F., Chamaillé-Jammes, S., Hebblewhite, M., Hopcraft, J. G. C., Merkle, J. A., Mueller, T., Mysterud, A., Peters, W., Roettger, C., Steingisser, A., Meacham, J. E., Abera, K., Adamczewski, J., Aikens, E. O., Bartlam-Brooks, H., Bennitt, E., Berger, J., Boyd, C., ... Zuther, S. (2021). Mapping out a future for ungulate migrations. *Science*, 372(6542), 566–569. https://doi. org/10.1126/science.abf0998
- Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye on life and planet. *Science*, *348*, aaa2478.
- Kays, R., Davidson, S. C., Berger, M., Bohrer, G., Fiedler, W., Flack, A., Hirt, J., Hahn, C., Gauggel, D., & Russell, B. (2022). The Movebank system for studying global animal movement and demography. *Methods in Ecology and Evolution*, 13, 419–431.
- King, A. J., Fehlmann, G., Biro, D., Ward, A. J., & Fürtbauer, I. (2018). Rewilding collective behaviour: An ecological perspective. *Trends in Ecology & Evolution*, 33, 347–357.
- Laskowski, K. L., Alirangues Nuñez, M. M., Hilt, S., Gessner, M. O., & Mehner, T. (2022). Predator group composition indirectly influences food web dynamics through predator growth rates. *The American Naturalist*, 199, 330–344.
- Lavelle, M. J., Hygnstrom, S. E., Hildreth, A. M., Campbell, T. A., Long, D. B., Hewitt, D. G., Beringer, J., & VerCauteren, K. C. (2012). Utility of improvised video-camera collars for collecting contact data from white-tailed deer: Possibilities in disease transmission studies. *Wildlife Society Bulletin*, 36, 828–834.
- Leibold, M. A., & Chase, J. M. (2017). *Metacommunity ecology*. Princeton University Press.
- Letten, A. D., Ke, P.-J., & Fukami, T. (2017). Linking modern coexistence theory and contemporary niche theory. *Ecological Monographs*, 87(2), 161–177. https://doi.org/10.1002/ecm.1242
- Levin, S. A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73, 1943–1967.
- Lowther, A. D., Lydersen, C., & Kovacs, K. M. (2015). A sum greater than its parts: Merging multi-predator tracking studies to increase ecological understanding. *Ecosphere*, 6, 1–13.
- MacArthur, R., & Levins, R. (1967). The limiting similarity, convergence, and divergence of coexisting species. *The American Naturalist*, 101, 377-385.
- Marvin, D. C., Koh, L. P., Lynam, A. J., Wich, S., Davies, A. B., Krishnamurthy, R., Stokes, E., Starkey, R., & Asner, G. P. (2016). Integrating technologies for scalable ecology and conservation. *Global Ecology and Conservation*, 7, 262–275.
- Merrick, M. J., & Koprowski, J. L. (2017). Should we consider individual behavior differences in applied wildlife conservation studies? *Biological Conservation*, 209, 34–44.
- Mertes, K., Jarzyna, M. A., & Jetz, W. (2020). Hierarchical multi-grain models improve descriptions of species' environmental associations, distribution, and abundance. *Ecological Applications*, 30(6), e02117.
- Milles, A., Dammhahn, M., & Grimm, V. (2020). Intraspecific trait variation in personality-related movement behavior promotes coexistence. *Oikos*, 129, 1441–1454.
- Milner, J. E., Blackwell, P. G., & Niu, M. (2021). Modelling and inference for the movement of interacting animals. *Methods in Ecology and Evolution*, 12, 54–69.
- Moll, R. J., Millspaugh, J. J., Beringer, J., Sartwell, J., & He, Z. (2007). A new 'view' of ecology and conservation through animal-borne video systems. *Trends in Ecology & Evolution*, 22, 660–668.
- Montgomery, R. A., Redilla, K. M., Ortiz-Calo, W., Smith, T., Keller, B., & Millspaugh, J. J. (2018). Evaluating the individuality of animalhabitat relationships. *Ecology and Evolution*, *8*, 10893–10901.
- Montiglio, P. O., Gotanda, K. M., Kratochwil, C. F., Laskowski, K. L., & Farine, D. R. (2019). Hierarchically embedded interaction networks

represent a missing link in the study of behavioral and community ecology. *Behavioral Ecology*, 31, 279–286.

- Murdoch, W. W., Briggs, C. J., & Nisbet, R. M. (2003). *Consumer-resource dynamics*. Princeton University Press.
- Nathan, R., Monk Christopher, T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., Baktoft, H., Beardsworth Christine, E., Bertram Michael, G., Bijleveld Allert, I., Brodin, T., Brooks Jill, L., Campos-Candela, A., Cooke Steven, J., Gjelland Karl, Ø., Gupte Pratik, R., Harel, R., Hellström, G., Jeltsch, F., ... Jarić, I. (2022). Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science*, 375, eabg1780.
- Noonan, M. J., Martinez-Garcia, R., Davis, G. H., Crofoot, M. C., Kays, R., Hirsch, B. T., Caillaud, D., Payne, E., Sih, A., Sinn, D. L., Spiegel, O., Fagan, W. F., Fleming, C. H., & Calabrese, J. M. (2021). Estimating encounter location distributions from animal tracking data. *Methods in Ecology and Evolution*, *12*, 1158–1173.
- Pansu, J., Guyton, J. A., Potter, A. B., Atkins, J. L., Daskin, J. H., Wursten, B., Kartzinel, T. R., & Pringle, R. M. (2019). Trophic ecology of large herbivores in a reassembling African ecosystem. *Journal of Ecology*, 107, 1355–1376.
- Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., & Araújo, M. B. (2011). *Ecological niches* and geographic distributions (MPB-49). Princeton University Press.
- Peterson, A. T., Soberón, J., Ramsey, J., & Osorio-Olvera, L. (2020). Cooccurrence networks do not support identification of biotic interactions. *Biodiversity Informatics*, 15, 1–10.
- Potts, J. R., Mokross, K., & Lewis, M. A. (2014). A unifying framework for quantifying the nature of animal interactions. *Journal of the Royal Society Interface*, 11, 20140333.
- Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters*, *3*, 349–361.
- Randin, C. F., Ashcroft, M. B., Bolliger, J., Cavender-Bares, J., Coops, N. C., Dullinger, S., Dirnböck, T., Eckert, S., Ellis, E., Fernández, N., Giuliani, G., Guisan, A., Jetz, W., Joost, S., Karger, D., Lembrechts, J., Lenoir, J., Luoto, M., Morin, X., ... Payne, D. (2020). Monitoring biodiversity in the Anthropocene using remote sensing in species distribution models. *Remote Sensing of Environment*, 239, 111626.
- Raymond, B., Lea, M. A., Patterson, T., Andrews-Goff, V., Sharples, R., Charrassin, J. B., Cottin, M., Emmerson, L., Gales, N., & Gales, R. (2015). Important marine habitat off East Antarctica revealed by two decades of multi-species predator tracking. *Ecography*, 38, 121–129.
- Schell, C. J., Stanton, L. A., Young, J. K., Angeloni, L. M., Lambert, J. E., Breck, S. W., & Murray, M. H. (2021). The evolutionary consequences of human-wildlife conflict in cities. *Evolutionary Applications*, 14, 178–197.
- Schirmer, A., Hoffmann, J., Eccard, J. A., & Dammhahn, M. (2020). My niche: Individual spatial niche specialization affects within- and between-species interactions. *Proceedings of the Royal Society B: Biological Sciences*, 287, 20192211.
- Schlägel, U. E., Grimm, V., Blaum, N., Colangeli, P., Dammhahn, M., Eccard, J. A., Hausmann, S. L., Herde, A., Hofer, H., Joshi, J., Kramer-Schadt, S., Litwin, M., Lozada-Gobilard, S. D., Müller, M. E. H., Müller, T., Nathan, R., Petermann, J. S., Pirhofer-Walzl, K., Radchuk, V., ... Jeltsch, F. (2020). Movement-mediated community assembly and coexistence. *Biological Reviews*, 95, 1073–1096.
- Schlägel, U. E., Signer, J., Herde, A., Eden, S., Jeltsch, F., Eccard, J. A., & Dammhahn, M. (2019). Estimating interactions between individuals from concurrent animal movements. *Methods in Ecology and Evolution*, 10, 1234–1245.
- Schmitz, O. J., Wilmers, C. C., Leroux, S. J., Doughty, C. E., Atwood, T. B., Galetti, M., Davies, A. B., & Goetz, S. J. (2018). Animals and the zoogeochemistry of the carbon cycle. *Science*, *362*, eaar3213.
- Silk, M. J., Finn, K. R., Porter, M. A., & Pinter-Wollman, N. (2018). Can multilayer networks advance animal behavior research? *Trends in Ecology & Evolution*, 33, 376–378.

- Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters*, 10, 1115–1123.
- Spiegel, O., Leu, S. T., Bull, C. M., & Sih, A. (2017). What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecology Letters*, 20, 3–18.
- Strandburg-Peshkin, A., Farine, D. R., Couzin, I. D., & Crofoot, M. C. (2015). Shared decision-making drives collective movement in wild baboons. *Science*, 348, 1358–1361.
- Strandburg-Peshkin, A., Farine, D. R., Crofoot, M. C., & Couzin, I. D. (2017). Habitat and social factors shape individual decisions and emergent group structure during baboon collective movement. *eLife*, *6*, e19505.
- Suraci, J. P., Smith, J. A., Chamaillé-Jammes, S., Gaynor, K. M., Jones, M., Luttbeg, B., Ritchie, E. G., Sheriff, M. J., & Sih, A. (2022). Beyond spatial overlap: Harnessing new technologies to resolve the complexities of predator-prey interactions. *Oikos*, e09004. https://doi. org/10.1111/oik.09004
- Thompson, P. L., Guzman, L. M., De Meester, L., Horváth, Z., Ptacnik, R., Vanschoenwinkel, B., Viana, D. S., & Chase, J. M. (2020). A processbased metacommunity framework linking local and regional scale community ecology. *Ecology Letters*, 23, 1314–1329.
- Tucker, M. A., Böhning-Gaese, K., Fagan, W. F., Fryxell, J. M., Van Moorter, B., Alberts, S. C., Ali, A. H., Allen, A. M., Attias, N., & Avgar, T. (2018). Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science*, 359, 466–469.
- Tuia, D., Kellenberger, B., Beery, S., Costelloe, B. R., Zuffi, S., Risse,
 B., Mathis, A., Mathis, M. W., van Langevelde, F., Burghardt,
 T., Kays, R., Klinck, H., Wikelski, M., Couzin, I. D., van Horn,
 G., Crofoot, M. C., Stewart, C. V., & Berger-Wolf, T. (2022).
 Perspectives in machine learning for wildlife conservation.
 Nature Communications, 13, 792.
- Urbano, F., Cagnacci, F., & Initiative, E. C. (2021). Data management and sharing for collaborative science: Lessons learnt from the Euromammals Initiative. *Frontiers in Ecology and Evolution*, *9*, 727023.
- Vanak, A. T., Fortin, D., Thaker, M., Ogden, M., Owen, C., Greatwood, S., & Slotow, R. (2013). Moving to stay in place: Behavioral mechanisms for coexistence of African large carnivores. *Ecology*, 94, 2619–2631.
- Veldhuis, M. P., Hofmeester, T. R., Balme, G., Druce, D. J., Pitman, R. T., & Cromsigt, J. P. G. M. (2020). Predation risk constrains herbivores' adaptive capacity to warming. *Nature Ecology & Evolution*, 4, 1069–1074.
- Villegas-Ríos, D., Freitas, C., Moland, E., Huneide Thorbjørnsen, S., & Olsen, E. M. (2020). Inferring individual fate from aquatic acoustic telemetry data. *Methods in Ecology and Evolution*, 11, 1186–1198.
- Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., Jung, V., & Messier, J. (2012). The return of the variance: Intraspecific variability in community ecology. *Trends in Ecology & Evolution*, 27, 244–252.
- Votier, S. C., Bearhop, S., Witt, M. J., Inger, R., Thompson, D., & Newton, J. (2010). Individual responses of seabirds to commercial fisheries revealed using GPS tracking, stable isotopes and vessel monitoring systems. *Journal of Applied Ecology*, 47, 487–497.
- Wikelski, M., Kays, R. W., Kasdin, N. J., Thorup, K., Smith, J. A., & Swenson, G. W. (2007). Going wild: What a global small-animal tracking system could do for experimental biologists. *Journal of Experimental Biology*, 210, 181–186.
- Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., de Grissac, S., Demšar, U., English, H. M., Franconi, N., Gómez-Laich, A., Griffiths, R. C., Kay, W. P., Morales, J. M., Potts, J. R., Rogerson, K. F., Rutz, C., Spelt, A., Trevail, A. M., Wilson, R. P., & Börger, L. (2020). Optimizing the use of biologgers for movement ecology research. *Journal of Animal Ecology*, *89*, 186–206.
- Wilmers, C. C., Nickel, B., Bryce, C. M., Smith, J. A., Wheat, R. E., & Yovovich, V. (2015). The golden age of bio-logging: How

animal-borne sensors are advancing the frontiers of ecology. *Ecology*, *96*, 1741–1753.

- Wilson, A. M., & Jetz, W. (2016). Remotely sensed high-resolution global cloud dynamics for predicting ecosystem and biodiversity distributions. *PLoS Biology*, 14, e1002415.
- Wootton, J. T. (1994). The nature and consequences of indirect effects in ecological communities. *Annual Review of Ecology and Systematics*, 25, 443–466.

How to cite this article: Costa-Pereira, R., Moll, R. J., Jesmer, B. R. & Jetz, W. (2022). Animal tracking moves community ecology: Opportunities and challenges. *Journal of Animal Ecology*, 91, 1334–1344. <u>https://doi.org/10.1111/1365-</u> 2656.13698